arXiv:math/0612060v1 [math.AG] 4 Dec 2006

NOTES ON THE CHERN-CHARACTER

HELGE MAAKESTAD

Abstract. Notes for some talks given at the seminar on characteristic classes at NTNU in autumn 2006. In the note a proof of the existence of a Chern-character from complex K-theory to any cohomology theory with values in graded **Q**-algebras equipped with a theory of characteristic classes is given. It respects the Adams and Steenrod operations.

Contents

1. Introduction 1
2. Euler classes and characteristic classes 2
3. Adams operations and Newton polynomials 4
4. The Chern-character and cohomology operations 8
References 12

1. Introduction

The aim of this note is to give an axiomatic and elementary treatment of Chern-characters of vectorbundles with values in a class of cohomology-theories arising in topology and algebra. Given a theory of Chern-classes for complex vectorbundles with values in singular cohomology one gets in a natural way a Chern-character from complex K-theory to singular cohomology using the projective bundle theorem and the Newton polynomials. The Chern-classes of a complex vectorbundle may be defined using the notion of an Euler class (see section 14 in [6]) and one may prove that a theory of Chern-classes with values in singular cohomology is unique. In this note it is shown one may relax the conditions on the theory for Chern-classes and still get a Chern-character. Hence the Chern-character depends on some choices.

Many cohomology theories which associate to a space a graded commutative **Q**-algebra H^* satisfy the projective bundle property for complex vectorbundles. This is true for De Rham-cohomology of a real compact manifold, singular cohomology of a compact topological space and complex K-theory. The main aim of this note is to give a self contained and elementary proof of the fact that any such cohomology theory will recieve a Chern-character from complex K-theory respecting the Adams and Steenrod operations.

Complex K-theory for a topological space B is considered, and characteristic classes in K-theory and operations on K-theory such as the Adams operations are constructed explicitly, following [5].

Date: November 2006.
Key words and phrases. Chern-character, Chern-classes, Euler classes, singular cohomology, De Rham-cohomology, complex K-theory, Adams operations, Steenrod operations.

The main result of the note is the following (Theorem 4.9):

Theorem 1.1. *Let* H^* *be any rational cohomology theory satisfying the projective bundle property. There is for all* $k \geq 1$ *a commutative diagram*

$$\begin{array}{ccc} K^*_{\mathbf{C}}(B) & \xrightarrow{Ch} & \mathrm{H}^{even}(B) \\ \downarrow{\scriptstyle \psi^k} & & \downarrow{\scriptstyle \psi^k_H} \\ K^*_{\mathbf{C}}(B) & \xrightarrow{Ch} & \mathrm{H}^{even}(B) \end{array}$$

where Ch is the Chern-character for H^**,* ψ^k *is the Adams operation and* ψ^k_H *is the Steenrod operation.*

The proof of the result is analogous to the proof of existence of the Chern-character for singular cohomology.

2. Euler classes and characteristic classes

In this section we consider axioms ensuring that any cohomology theory H^* satisfying these axioms, recieve a Chern-character for complex vectorbundles. By a cohomology theory we mean a contravariant functor

$$\mathrm{H}^* : \underline{Top} \to \underline{\mathbf{Q} - algebras}$$

from the category of topological spaces to the category of graded commutative $\mathbf{Q}$-algebras with respect to continuous maps of topological spaces. We say the theory satisfy the *projective bundle property* if the following axioms are satisfied: For any rank n complex continuous vectorbundle E over a compact space B There is an *Euler class*

$$u_E \in \mathrm{H}^2(\mathbf{P}(E)) \tag{2.0.1}$$

where $\pi : \mathbf{P}(E) \to B$ is the projective bundle associated to E. This assignment satisfy the following properties: The Euler class is natural, i.e for any map of topological spaces $f : B' \to B$ it follows

$$f^* u_E = u_{f^* E} \tag{2.0.2}$$

For $E = \oplus_{i=1}^n L_i$ where L_i are linebundles there is an equation

$$\prod_{i=1}^{n} (u_E - \pi^* u_{L_i}) = 0 \text{ in } \mathrm{H}^{2n}(\mathbf{P}(E)) \tag{2.0.3}$$

The map π^* induce an injection $\pi^* : \mathrm{H}^*(B) \to \mathrm{H}^*(\mathbf{P}(E))$ and there is an equality

$$\mathrm{H}^*(\mathbf{P}(E)) = \mathrm{H}^*(B)\{1, u_E, u_E^2, .., u_E^{n-1}\}.$$

Assume H^* satisfy the projective bundle property. There is by definition an equation

$$u_E^n - c_1(E) u_E^{n-1} + \cdots + (-1)^n c_n(E) = 0$$

in $\mathrm{H}^*(\mathbf{P}(E))$.

Definition 2.1. The class $c_i(E) \in \mathrm{H}^{2i}(B)$ is the *i'th characteristic class* of E.

Example 2.2. *If $\mathbf{P}(E) \to B$ is the projective bundle of a complex vector bundle and $u_E = e(\lambda(E)) \in \mathrm{H}^2(\mathbf{P}(E), \mathbf{Z})$ is the Euler classe of the tautological linebundle $\lambda(E)$ on $\mathbf{P}(E)$ in singular cohomology as defined in Section 14* [6]*, one verifies the properties above are satisfied. One gets the Chern-classes $c_i(E) \in \mathrm{H}^{2i}(B, \mathbf{Z})$ in singular cohomology.*

Definition 2.3. A *theory of characteristic classes* with values in a cohomology theory H^* is an assignment

$$E \to c_i(E) \in \mathrm{H}^{2i}(B)$$

for every complex finite rank vectorbundle E on B satisfying the following axioms:

$$f^* c_i(E) = c_i(f^* E) \tag{2.3.1}$$

$$\text{If } E \cong F \text{ it follows } c_i(E) = c_i(F) \tag{2.3.2}$$

$$c_k(E \oplus F) = \sum_{i+j=k} c_i(E) c_j(F). \tag{2.3.3}$$

Note: if $\phi : \mathrm{H}^* \to \mathrm{H}^*$ is a functorial endomorphism of H^* which is a ring-homomorphism and c is a theory of characteristic classes, it follows the assignment $E \to \overline{c}_i(E) = \phi(c_i(E))$ is a theory of characteristic classes.

Example 2.4. *Let $k \in \mathbf{Z}$ and let ψ_H^k be the ring-endomorphism of H^{even} defined by $\psi_H^k(x) = k^r x$ where $x \in \mathrm{H}^{2r}(B)$. Given a theory $c_i(E)$ satisfying Definition 2.3 it follows $\overline{c}_i(E) = \psi_H^k(c_i(E))$ is a theory satisfying Definition 2.3.*

Note furthermore: Assume γ_1 is the tautological linebundle on $\mathbf{P}^1$. Since we do not assume $c_1(\gamma_1) = z$ where z is the canonical generator of $\mathrm{H}^2(\mathbf{P}^1, \mathbf{Z})$ it does not follow that an assignment $E \to c_i(E)$ is uniquely determined by the axioms $2.3.1 - 2.3.3$. We shall see later that the axioms $2.3.1 - 2.3.3$ is enough to define a Chern-character.

Theorem 2.5. *Assume the theory H^* satisfy the projective bundle property. It follows H^* has a theory of characteristic classes.*

Proof. We verify the axioms for a theory of characteristic classes. Axiom 2.3.1: Assume we have a map of rank n bundles $f : F \to E$ over a map of topological spaces $g : B' \to B$. We pull back the equation

$$u_E^n - c_1(E) u_E^{n-1} + \cdots + (-1)^n c_n(E) = 0$$

in $\mathrm{H}^{2n}(\mathbf{P}(E))$ to get an equation

$$u_F^n - f^* c_1(E) u_F^{n-1} + \cdots + (-1)^n f^* c_n(E) = 0$$

and by unicity we get $f^* c_i(E) = c_i(F)$. It follows $c_i(E) = c_i(F)$ for isomorphic bundles E and F, hence Axiom 2.3.2 is ok. Axiom 2.3.3: Assume $E \cong \oplus_{i=1}^n L_i$ is a decomposition into linebundles. There is an equation $\prod(u_E - u_{L_i})$ hence we get a polynomial relation

$$u_E^n - s_1(u_{L_i}) u_E^{n-1} + \cdots + (-1)^n s_n(u_{L_i}) = 0$$

in $\mathrm{H}^{2n}(\mathbf{P}(E))$. Since $c_1(L_i) = -u_{L_i}$ it follows

$$\prod(c(L_i)) = \prod(1 + c_1(L_i)) = c(E)$$

and this is ok. □

Given a compact topological space B. We may consider the Grothendieck-ring $K^*_{\mathbf{C}}(B)$ of complex finite-dimensional vectorbundles. It is defined as the free abelian group on isomorphism-classes $[E]$ where E is a complex vectorbundle, modulo the subgroup generated by elements of the type $[E \oplus F] - [E] - [F]$. It has direct sum as additive operation and tensor product as multiplication. Assume E is a complex vectorbundle of rank n and let

$$\pi : \mathbf{P}(E) \to B$$

be the associated projective bundle. We have a *projective bundle theorem* for complex K-theory:

Theorem 2.6. *The group* $\mathrm{K}^*(\mathbf{P}(E))$ *is a free* $\mathrm{K}^*(B)$ *module of finite rank with generator* u *- the euler class of the tautological line-bundle. The elements* $\{1, u, u^2, .., u^{n-1}\}$ *is a free basis.*

Proof. See Theorem IV.2.16 in [5]. □

As in the case of singular cohomology, we may define characteristic classes for complex bundles with values in complex K-theory using the projective bundle theorem: The element u^n satisfies an equation

$$u^n - c_1(E)u^{n-1} + c_2(E)u^{n-2} + \cdots + (-1)^{n-1}c_{n-1}(E)u + (-1)^n c_n(E) = 0$$

in $\mathrm{K}^*(\mathbf{P}(E))$. One verifies the axioms defined above are satisfied, hence one gets characteristic classes $c_i(E) \in \mathrm{K}^*_{\mathbf{C}}(B)$ for all $i = 0, .., n$

Theorem 2.7. *The characteristic classes* $c_i(E)$ *satisfy the following properties:*

$$f^* c_i(E) = c_i(f^* E) \tag{2.7.1}$$

$$c_k(E \oplus F) = \sum_{i+j=k} c_i(E)c_j(F) \tag{2.7.2}$$

$$c_1(L) = 1 - L \qquad\qquad c_i(L) = 0, i > 1 \tag{2.7.3}$$

where E *is any vectorbundle, and* L *is a line bundle.*

Proof. See Theorem IV.2.17 in [5]. □

3. Adams operations and Newton polynomials

We introduce some cohomology operations in complex K-theory and Newton-polynomials and prove elementary properties following the book [5].

Let $\Phi(B)$ be the abelian monoid of elements of the type $\sum n_i[E_i]$ with $n_i \geq 0$. Consider the bundle $\lambda^i(E) = \wedge^i E$ and the association

$$\lambda_t(E) = \sum_{i \geq 0} \lambda^i(E) t^i$$

giving a map

$$\lambda_t = \Phi(X) \to 1 + t\, \mathrm{K}^*_{\mathbf{C}}(B)[[t]]$$

One checks

$$\lambda_t(E \oplus F) = \lambda_t(E)\lambda_t(F)$$

hence the map λ_t is a map of abelian monoids, hence gives rise to a map

$$\lambda_t : \mathrm{K}^*_{\mathbf{C}}(B) \to 1 + t\, \mathrm{K}^*_{\mathbf{C}}(B)[[t]]$$

from the additive abelian group $\mathrm{K}^*_{\mathbf{C}}(B)$ to the set of powerseries with constant term equal to one. Explicitly the map is as follows:

$$\lambda_t(n[E] - m[F]) = \lambda_t(E)^n \lambda_t(F)^{-m}.$$

When n denotes the trivial bundle of rank n we get the explicit formula

$$\lambda_t([E] - n) = \lambda_t(E)(1+t)^{-n}.$$

Let $u = t/1 - t$. We may define the new powerseries

$$\gamma_t(E) = \lambda_u(E) = \sum_{k \geq 0} \lambda^i(E) u^i.$$

It follows

$$\gamma_t(E \oplus F) = \lambda_u(E \oplus F) = \lambda_u(E)\lambda_u(F) = \gamma_t(E)\gamma_t(F).$$

We may write formally

$$\gamma_t(E) = \sum_{k \geq 0} \gamma^i(E) t^i \in K^*_{\mathbf{C}}(B)[[t]]$$

hence it follows that

$$\gamma^k(E) = \sum_{i+j=k} \gamma^i(E)\gamma^j(E).$$

We get operations

$$\gamma^i : \mathrm{K}^*_{\mathbf{C}}(B) \to \mathrm{K}^*_{\mathbf{C}}(B)$$

for all $i \geq 1$. We next define *Newton polynomials* using the elementary symmetric functions. Let $u_1, u_2, u_3, ..$ be independent variables over the integers $\mathbf{Z}$, and let $Q_k = u_1^k + u_2^k + \cdots + u_k^k$ for $k \geq 1$. It follows Q_k is invariant under permutations of the variables u_i: for any $\sigma \in S_k$ we have $\sigma Q_k = Q_k$ hence we may express Q_k as a polynomial in the elementary symmetric functions σ_i:

$$Q_k = Q_k(\sigma_1, \sigma_2, .., \sigma_k).$$

We define

$$s_k(\sigma) = Q_k(\sigma_1, \sigma_2, .., \sigma_k)$$

to be the $k'th$ Newton polynomial in the variables $\sigma_1, \sigma_2, .., \sigma_k$ where σ_i is the $i'th$ elementary symmetric function. One checks the following:

$$s_1(\sigma_1) = \sigma_1,$$

$$s_2(\sigma_1, \sigma_2) = \sigma_1^2 - 2\sigma_2,$$

and

$$s_2(\sigma_1, \sigma_2, \sigma_3) = \sigma_1^3 - 3\sigma_1\sigma_2 + 3\sigma_3$$

and so on.

Let $n \geq 1$ and consider the polynomial

$$p(1) = (1 + tu_1)(1 + tu_2) \cdots (1 + tu_n) = t^n\sigma_n + t^{n-1}\sigma_{n-1} + \cdots + t\sigma_1 + 1$$

where

$$\sigma_i = \sigma_i(u_1, .., u_n)$$

is the ith elementary symmetric polynomial in the variables $u_1, u_2, .., u_n$.

Lemma 3.1. *There is an equality*

$$Q_k(\sigma_1(u_1, .., u_n), \sigma_2(u_1, .., u_n), .., \sigma_k(u_1, .., u_n)) = u_1^k + u_2^k + \cdots + u_n^k.$$

Proof. Trivial. □

Assume we have virtual elements $x = E - n = \oplus^n(L_i - 1)$ and $y = F - p = \oplus^p(R_j - 1)$ in complex K-theory $\mathrm{K}^*_{\mathbf{C}}(B)$. We seek to define a cohomology-operation c on complex K-theory using a formal powerseries

$$f(u) = a_1 u + a_2 u^2 + a_3 u^3 + \cdots \in \mathbf{Z}[[u]].$$

We define the element

$$c(x) = a_1 Q_1(\gamma^1(x)) + a_2 Q_2(\gamma^1(x), \gamma^2(x)) + a_3 Q_3(\gamma^1(x), \gamma^2(x), \gamma^3(x)) + \cdots$$

Proposition 3.2. *Let L be a linebundle. Then $\gamma_t(L-1) = 1+t(L-1) = 1-c_1(L)t$. Hence $\gamma^1(L-1) = L-1$ and $\gamma^i(L-1) = 0$ for $i > 1$.*

Proof. We have by definition

$$\gamma_t(E) = \lambda_u(E) = \sum_{k \geq 0} \lambda^k(E) u^k = \sum_{k \geq 0} \lambda^k(E)(t/1-t)^k.$$

We have that

$$\gamma_t(nE - mF) = \lambda_u(E)^n \lambda_u(F)^{-m}.$$

We get

$$\gamma_t(L-1) = \lambda_u(L)\lambda_u(1)^{-1}.$$

We have

$$\lambda_t(n) = (1+t)^n$$

hence

$$\gamma_t(n) = \lambda_u(n) = (1+u)^n = (1 + t/1-t)^n = (1-t)^{-n}.$$

We get:

$$\gamma_t(L-1) = \gamma_t(L)\gamma_t(1)^{-1} = \lambda_u(L)(1-t)^{-1} =$$
$$(1+Lu)(1-t)^{-1} = (1 + L(t/t-1))(1-t)^{-1} =$$
$$\frac{1+t(L-1)}{1-t}(1-t) = 1 + t(L-1) = 1 - c_1(L)t.$$

And the proposition follows. □

Note: if $x = L - 1$ we get

$$c(x) = \sum_{k \geq 0} a_k Q_k(\gamma^1(x), \gamma^2(x), .., \gamma^k(x)) =$$
$$\sum_{k \geq 1} a_k Q_k(\gamma^1(x), 0, ..., 0) = \sum_{k \geq 1} a_k \gamma^1(x)^k =$$
$$\sum_{k \geq 1} a_k (L-1)^k = \sum_{k \geq 0} (-1)^k a_k c_1(L)^k.$$

We state a Theorem:

Theorem 3.3. *Let $E \to B$ be a complex vectorbundle on a compact topological space B. There is a map $\pi : B' \to B$ such that $\pi^* E$ decompose into linebundles, and the map $\pi^* : \mathrm{H}^*(B) \to \mathrm{H}^*(B')$ is injective.*

Proof. See [5] Theorem IV.2.15. □

Note: By [5] Proposition II.1.29 there is a split exact sequence

$$0 \to K'_{\mathbf{C}}(B) \to \mathrm{K}^*_{\mathbf{C}}(B) \to \mathrm{H}^0(B, \mathbf{Z}) \to 0$$

hence the group $K'_{\mathbf{C}}(B)$ is generated by elements of the form $E - n$ where E is a rank n complex vectorbundle.

Proposition 3.4. *The operation c is additive, i.e for any* $x, y \in \mathrm{K}^*_{\mathbf{C}}(B)$ *we have*

$$c(x+y) = c(x) + c(y).$$

Proof. The proof follows the proof in [5], Proposition IV.7.11. We may by the remark above assume $x = E - n$ and $y = F - p$ where $x, y \in K'_{\mathbf{C}}(B)$. We may also from Theorem 3.3 assume $E = \oplus^n L_i$ and $F = \oplus^p R_j$ where L_i, R_j are linebundles. We get the following:

$$\gamma_t(x+y) = \prod^n \gamma_t(L_i - 1) \prod^p \gamma_t(R_j - 1) = \prod^n (1 + tu_i) \prod^p (1 + tv_j) =$$

$$t^{n+p}\sigma_{n+p}(u_1, .., u_n, v_1, .., v_p) + t^{n+p-1}\sigma_{n+p-1}(u_1, .., u_n, v_1, .., v_p) +$$

$$\cdots + t\sigma_1(u_1, .., u_n, v_1, .., v_p) + 1$$

hence

$$\gamma^i(x+y) = \sigma_i(u_1, .., u_n, v_1, .., v_p).$$

We get:

$$Q_k(\gamma^1(x+y), .., \gamma^k(x+y)) = Q_k(\sigma_1(u_i, v_j), .., \sigma_k(u_i, v_j))$$

which by Lemma 3.1 equals

$$u_1^k + \cdots u_n^k + v_1^k + \cdots v_p^k = Q_k(\sigma_1(u_i), .., \sigma_k(u_i)) + Q_k(\sigma_1(v_j), .., \sigma_k(v_j)) =$$

$$Q_k(\gamma^i(x)) + Q_k(\gamma^i(y)).$$

We get:

$$c(x+y) = \sum_{k \geq 0} a_k Q_k(\gamma^i(x+y)) =$$

$$\sum_{k \geq 0} a_k Q_k(\gamma^i(x)) + \sum_{k \geq 0} a_k Q_k(\gamma^i(y)) = c(x) + c(y)$$

and the claim follows.

□

We may give an explicit and elementary construction of the Adams-operations:

Theorem 3.5. *Let* $k \geq 1$. *There are functorial operations*

$$\psi^k : \mathrm{K}^*_{\mathbf{C}}(B) \to \mathrm{K}^*_{\mathbf{C}}(B)$$

with the properties

$$\psi^k(x+y) = \psi^k(x) + \psi^k(y) \tag{3.5.1}$$

$$\psi^k(L) = L^k \tag{3.5.2}$$

$$\psi^k(xy) = \psi^k(x)\psi^k(y) \tag{3.5.3}$$

$$\psi^k(1) = 1 \tag{3.5.4}$$

where L is a line bundle. The operations ψ^k *are the only operations that are ring-homomorphisms - the* Adams operations

Proof. We need:

$$\psi^k(L-1) = \psi^k(L) - \psi^k(1) = L^k - 1.$$

We have in K-theory:

$$L^k - 1 = (L - 1 + 1)^k - 1 = \sum_{i \geq 0} \binom{k}{i} (L-1)^{k-i} 1^i - 1 =$$

$$\binom{k}{1}(L-1)+\binom{k}{2}(L-1)^2+\cdots+\binom{k}{k}(L-1)^k.$$

We get the series

$$c=\sum_{i=1}^{k}\binom{k}{i}u^k\in\mathbf{Z}[[u]].$$

The following operator

$$\psi^k=\sum_{i=1}^{k}\binom{k}{i}Q_i(\gamma^1,...,\gamma^i)$$

is an explicit construction of the Adams-operator. One may verify the properties in the theorem, and the claim follows. □

Assume E, F are complex vectorbundles on B and consider the Chern-polynomial

$$c_t(E\oplus F)=1+c_1(E\oplus F)t+\cdots+c_N(E\oplus F)t^N$$

where $N=rk(E)+rk(F)$. Assume there is a decomposition $E=\oplus^n L_i$ and $F=\oplus^p R_j$ into linebundles. We get a decomposition

$$c_t(E\oplus F)=\prod c_t(L_i)\prod c_t(R_j)=(1+a_1t)(1+a_2t)\cdots(1+b_1t)\cdots(1+b_pt)$$

where $a_i=c_1(L_i), b_j=c_1(R_j)$. We get thus

$$c_i(E\oplus F)=\sigma_i(a_1,..,a_n,b_1,..,b_p).$$

Let

$$Q_k=u_1^k+\cdots+u_k^k=Q_k(\sigma_1,..,\sigma_k)$$

where σ_i is the ith elementary symmetric function in the u_i's.

Proposition 3.6. *The following holds:*

$$Q_k(c_1(E\oplus F),..,c_k(E\oplus F))=Q_K(c_i(E))+Q_k(c_i(F)).$$

Proof. We have

$$Q_k(c_i(E\oplus F))=Q_k(\sigma_i(a_i,b_j))=$$
$$a_1^k+\cdots a_n^k+b_1^k+\cdots b_p^k=Q_k(c_i(E))+Q_k(c_i(F))$$

and the claim follows. □

4. The Chern-character and cohomology operations

We construct a Chern-character with values in singular cohomology, using Newton-polynomials and characteristic classes following [5]. The $k'th$ Newton-classe $s_k(E)$ of a complex vectorbundle will be defined using characteristic classes of E: $c_1(E),..,c_k(E)$ and the $k'th$ Newton-polynomial $s_k(\sigma_1,..,\sigma_k)$. We us this construction to define the Chern-character $Ch(E)$ of the vectorbundle E.

We first define *Newton polynomials* using the elementary symmetric functions. Let $u_1,u_2,u_3,..$ be independent variables over the integers $\mathbf{Z}$, and let $Q_k=u_1^k+u_2^k+\cdots+u_k^k$ for $k\geq 1$. It follows Q_k is invariant under permutations of the variables u_i: for any $\sigma\in S_k$ we have $\sigma Q_k=Q_k$ hence we may express Q_k as a polynomial in the elementary symmetric functions σ_i:

$$Q_k=Q_k(\sigma_1,\sigma_2,..,\sigma_k).$$

We define

$$s_k(\sigma)=Q_k(\sigma_1,\sigma_2,..,\sigma_k)$$

to be the $k'th$ Newton polynomial in the variables $\sigma_1, \sigma_2, .., \sigma_k$ where σ_i is the $i'th$ elementary symmetric function. One checks the following:

$$s_1(\sigma_1) = \sigma_1,$$

$$s_2(\sigma_1, \sigma_2) = \sigma_1^2 - 2\sigma_2,$$

and

$$s_2(\sigma_1, \sigma_2, \sigma_3) = \sigma_1^3 - 3\sigma_1\sigma_2 + 3\sigma_3$$

and so on.

Assume we have a cohomology theory H^* satisfying the projective bundle property. One gets characteristic classes $c_i(E)$ for a complex vectorbundle E on B:

$$c_i(E) \in \mathrm{H}^{2i}(B).$$

Let the class $S_k(E) = s_k(c_1(E), c_2(E), .., c_k(E)) \in \mathrm{H}^{2k}(B)$ be the $k'th$ Newton-class of the bundle E. One gets:

$$s_k(\sigma_1, 0, .., 0) = \sigma_1^k$$

for all $k \geq 1$. Assume E, F linebundles. We see that

$$S_2(E \oplus F) = c_1(E \oplus F)^2 - 2c_2(E \oplus F) =$$

$$(c_1(E) + c_1(F))^2 - 2(c_2(E) + c_1(E)c_1(F) + c_2(F)) =$$

$$c_1(E)^2 + 2c_1(E)c_1(F) + c_1(F)^2 - 2c_2(E) - 2c_1(E)c_1(F) - 2c_2(F) =$$

$$c_1(E)^2 - 2c_2(E) + c_1(F)^2 - 2c_2(F) = S_2(E) + S_2(F).$$

This holds in general:

Proposition 4.1. *For any vectorbundles E, F we have the formula*

$$S_k(E \oplus F) = S_k(E) + S_k(F).$$

Proof. This follows from 3.6. □

Let $\mathrm{K}^*_{\mathbf{C}}(B)$ be the Grothendieck-group of complex vectorbundles on B, i.e the free abelian group modulo exact sequences $K^*_{\mathbf{C}}(B) = \oplus\mathbf{Z}[E]/U$ where U is the subgroup generated by elements $[E \oplus F] - [E] - [F]$.

Definition 4.2. The class

$$Ch(E) = \sum_{k \geq 0} \frac{1}{k!} S_k(E) \in \mathrm{H}^{even}(B)$$

is the *Chern-character* of E.

Lemma 4.3. *The Chern-character defines a group-homomorphism*

$$Ch : \mathrm{K}^*_{\mathbf{C}}(B) \to \mathrm{H}^{even}(B)$$

between the Grothendieck group $\mathrm{K}^*_{\mathbf{C}}(B)$ and the even cohomology of B with rational coefficients.

Proof. By Proposition 4.1 we get the following: For any E, F we have

$$Ch(E \oplus F) = \sum_{k \geq 0} \frac{1}{k!} s_k(E \oplus F) = \sum_{k \geq 0} \frac{1}{k!}(s_k(E) + s_k(F)) =$$

$$\sum_{k \geq 0} \frac{1}{k!} s_k(E) + \sum_{k \geq 0} \frac{1}{k!} s_k(F) = Ch(E) + Ch(F).$$

We get

$$Ch([E \oplus F] - [E] - [F]) = Ch(E \oplus F) - Ch(E) - Ch(F) = 0$$

and the Lemma follows. □

Example 4.4. *Given a real continuous vectorbundle F on B there exist Stiefel-Whitney classes $w_i(F) \in \mathrm{H}^i(B, \mathbf{Z}/2)$ (see [6]) satisfying the necessary conditions, and we may define a "Chern-character"*

$$Ch : \mathrm{K}^*_{\mathbf{R}}(B) \to \mathrm{H}^*(B, \mathbf{Z}/2)$$

by

$$Ch(F) = \sum_{k \geq 0} Q_k(w_1(F), .., w_k(F)).$$

This gives a well-defined homomorphism of abelian groups because of the universal properties of the Newton-polynomials and the fact $\mathrm{H}^*(B, \mathbf{Z}/2)$ is commutative. The formal properties of the Stiefel-Whitney classes w_i ensures that for real bundles E, F Proposition 3.6 still holds: We have the formula

$$Q_k(w_i(E \oplus F)) = Q_k(w_i(E)) + Q_k(w_i(F)).$$

Since $S_k(\sigma_1, 0, ..., 0) = \sigma_1^k$ we get the following: When E, F are linebundles we have:

$$S_k(E \otimes F) = S_k(c_1(E \otimes F), 0, .., 0) = (c_1(E \otimes F))^k = (c_1(E) + c_1(F))^k =$$

$$\sum_{i+j=k} \binom{i+j}{i} c_1(E)^i c_1(F)^j = \sum_{i+j=k} \binom{i+j}{i} S_i(E) S_j(F).$$

This property holds for general E, F:

Proposition 4.5. *Let E, F be complex vectorbundles on a compact topological space B. Then the following formulas hold:*

$$S_k(E \otimes F) = \sum_{i+j=k} \binom{i+j}{i} S_i(E) S_j(E) \tag{4.5.1}$$

Proof. We prove this using the splitting-principle and Proposition 4.1. Assume E, F are complex vectorbundles on B and $f : B' \to B$ is a map of topological spaces such that $f^*E = \oplus_i L_i$, $f^*F = \oplus_j M_j$ where L_i, M_j are linebundles and the pull-back map $f^* : \mathrm{H}^*(B) \to \mathrm{H}^*(B')$ is injective. We get the following calculation:

$$f^* S_k(E \otimes F) = S_k(f^*E \otimes F) = S_k(\oplus L_i \otimes M_j)$$

hence by Lemma 4.1 we get

$$\sum_{i,j} S_k(L_i \otimes M_j) = \sum_i (\sum_j S_k(L_i \otimes M_j)) =$$

$$\sum_i \sum_j \sum_{u+v=k} \binom{u+v}{u} S_u(L_i) S_v(M_j) =$$

$$\sum_i \sum_{u+v=k} \binom{u+v}{u} S_u(L_i) S_v(\oplus M_j) =$$

$$\sum_{u+v=k} \binom{u+v}{u} S_u(\oplus L_i) S_v(\oplus M_j) =$$

$$\sum_{u+v=k}\binom{u+v}{u}S_u(f^*E)S_v(f^*F) = f^*\sum_{u+v=k}\binom{u+v}{u}S_u(E)S_v(F),$$

and the result follows since f^* is injective. □

Theorem 4.6. *The Chern-character defines a ring-homomorphism*

$$Ch : K^*_{\mathbf{C}}(B) \to \mathrm{H}^{even}(B).$$

Proof. From Proposition 4.5 we get:

$$Ch(E\otimes F) = \sum_{k\geq 0}\frac{1}{k!}S_k(E\otimes F) =$$

$$\sum_{k\geq 0}\frac{1}{k!}\sum_{i+j=k}\binom{i+j}{i}S_i(E)S_j(F) =$$

$$(\sum_{k\geq 0}\frac{1}{k!}S_k(E))(\sum_{k\geq 0}\frac{1}{k!}S_k(F) = Ch(E)Ch(F)$$

and the Theorem is proved. □

Example 4.7. *For complex K-theory* $\mathrm{K}^*_{\mathbf{C}}(B)$ *we have for any complex vectorbundle* E *characteristic classes* $c_i(E) \in \mathrm{K}^*_{\mathbf{C}}(B)$ *satisfying the neccessary conditions, hence we get a group-homomorphism*

$$Ch_{\mathbf{Z}} : \mathrm{K}^*_{\mathbf{C}}(B) \to \mathrm{K}^*_{\mathbf{C}}(B)$$

defined by

$$Ch_{\mathbf{Z}}(E) = \sum_{k\geq 0} Q_k(c_1(E),..,c_k(E)).$$

If we tensor with the rationals, we get a ring-homomorphism

$$Ch_{\mathbf{Q}} : \mathrm{K}^*_{\mathbf{C}}(B) \to \mathrm{K}^*_{\mathbf{C}}(B)\otimes\mathbf{Q}$$

defined by

$$Ch(E) = \sum_{k\geq 0}\frac{1}{k!}Q_k(c_1(E),..,c_k(E)).$$

Theorem 4.8. *Let* B *be a compact topological space. The Chern-character*

$$Ch^{\mathbf{Q}} : K^*_{\mathbf{C}}(B)\otimes\mathbf{Q} \to \mathrm{H}^{even}(B,\mathbf{Q})$$

is an isomorphism. Here $\mathrm{H}^*(B,\mathbf{Q})$ *denotes singular cohomology with rational coefficients.*

Proof. See [5]. □

The Chern-character is related to the Adams-operations in the following sense: There is a ring-homomorphism

$$\psi^k_H : \mathrm{H}^{even}(B) \to \mathrm{H}^{even}(B)$$

defined by

$$\psi^k_H(x) = k^r x$$

when $x \in \mathrm{H}^{2r}(B)$. The Chern-character respects these cohomology operations in the following sense:

Theorem 4.9. *There is for all $k \geq 1$ a commutative diagram*

$$\begin{array}{ccc} K^*_{\mathbf{C}}(B) & \xrightarrow{Ch} & \mathrm{H}^{even}(B) \\ \downarrow{\scriptstyle \psi^k} & & \downarrow{\scriptstyle \psi^k_H} \\ K^*_{\mathbf{C}}(B) & \xrightarrow{Ch} & \mathrm{H}^{even}(B) \end{array}$$

where ψ^k is the Adams operation defined in the previous section.

Proof. The proof follows Theorem V.3.27 in [5]: We may assume L is a linebundle and we get the following calculation: $\psi^k(L) = L^k$ and $c_1(L^k) = kc_1(L)$ hence

$$Ch(\psi^k(L)) = exp(kc_1(L)) = \sum_{i \geq 0} \frac{1}{i!} k^i c_1(L)^i =$$

$$\psi^k_H(exp(c_1(L))) = \psi^k_H(Ch(L)$$

and the claim follows. □

Hence the Chern-character is a morphism of cohomology-theories respecting the additional structure given by the Adams and Steenrod-operations.

References

[1] J. Dupont, Curvature and characteristic classes, Lecture Notes in Mathematics Vol. 640 *Springer Verlag* (1978)
[2] W. Fulton, S. Lang, Riemann-Roch algebra, *Grundlehren Math. Wiss.* no. 277 (1985)
[3] A. Grothendieck, Theorie des classes de Chern, *Bull. Soc. Math. France* no. 86 (1958)
[4] D. Husemoeller, Fibre bundles, GTM (1979)
[5] M. Karoubi, K-theory - an introduction, *Grundlehren Math. Wiss.* (1978)
[6] J. Milnor, Characteristic Classes (1966)
[7] N. Steenrod, Cohomology operations, *Princeton University Press* (1962)
[8] W. End, Über Adams-operationen, *Invent. Math* no. 9 (1969)

NTNU, Trondheim
E-mail address: Helge.Maakestad@math.ntnu.no